### CLINICAL VIDEO



# Ataxic gait

## | Tengku Ahmad Hidayat Tengku Kamadjaja Aziz<sup>2</sup> | Shalini Bhaskar<sup>1,2</sup> Vanita Paramajothy<sup>2</sup>

#### Correspondence

Shalini Bhaskar, Department of Medicine, Universiti Teknologi MARA, Shah Alam,

Email: drshalini@uitm.edu.my

## Abstract

Ataxia as the main complaint in a young patient prompts to diagnose neurodegenerative diseases. Not all ataxias are irreversible. We present a video of a patient in the clinic with ataxia diagnosed as subacute combined degeneration of the cord.



FIGURE 1 Sagittal T2-weighted images reveal an intraspinal hyperintensity of the cervical and thoracic spinal cord with no mass effect

The young man presented with unsteadiness, lethargy, and tingling for 3 months. He was pale with cognitive impairment (Montreal Cognitive Assessment score 25/30). He underscored in repeating the digits backward, serial seven subtraction, and delayed recall. He had ataxic gait, increased deep tendon reflexes, bilateral positive Babinski, impaired vibration and joint position sense, and positive Romberg's sign (Video S1). He had macrocytic anemia, low serum B12, positive intrinsic factor, and anti-parietal antibodies. MRI is described in Figure 1. Therefore, sensory ataxia due to subacute combined degeneration of the cord and pernicious anemia was diagnosed. He improved with B12 therapy. Video S2.

#### **ACKNOWLEDGMENTS**

We wish to thank the Medical director of Thomson Hospital Kota Damansara for granting permission to report this case.

## CONFLICT OF INTEREST

None declared.

## **AUTHOR CONTRIBUTIONS**

SB wrote the draft of the paper. TAHTKA helped in the management of the patient. The images were contributed by VP. All authors reviewed and edited the paper and approved the final version of the paper.

This is an open access article under the terms of the Creative Commons Attribution NonCommercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non commercial and no modifications or adaptations are made. © 2021 The Authors. Clinical Case Reports published by John Wiley & Sons Ltd.

<sup>&</sup>lt;sup>1</sup>Department of Medicine, Universiti Teknologi MARA, Shah Alam, Malaysia

<sup>&</sup>lt;sup>2</sup>Department of Medicine, Thomson Hospital Kota Damansara, Petaling Jaya, Malaysia

## ETHICAL APPROVAL

Ethics approval was waivered by the Thomson Hospital Kota Damansara Ethics Committee due to the nature of the manuscript (ie, case report). The manuscript DOES NOT report on any animal data or tissue.

#### INFORMED CONSENT

Written informed consent was obtained from the patient for their information to be published in this article.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from Thomson Hospital Kota Damansara, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of Thomson Hospital Kota Damansara.

#### ORCID

Shalini Bhaskar https://orcid.org/0000-0002-9169-4209

#### REFERENCE

 Langan R, Goodbred A. Vitamin B12 deficiency: recognition and management. 2020. https://www.aafp.org/afp/2017/0915/p384. html. Accessed August 5, 2020

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Bhaskar S, Tengku Kamadjaja Aziz TAH, Paramajothy V. Ataxic gait. *Clin Case Rep.* 2021;9:e04388. https://doi.org/10.1002/ccr3.4388